

Department of Ophthalmology Santa Rosa

Name:	
MR#:	
	Imprint Area

# POSTOPERATIVE INSTRUCTIONS: ENUCLEATION

#### WHAT YOU MAY EXPECT IN THE FIRST HOURS AFTER SURGERY

Sore Socket

Sleepiness

Poor Appetite

Possible Nausea and Vomiting (first 1-8 hours)

Do not remove the bandage

#### WHAT TO TAKE FOR PAIN

1. For pain relief, take Tylenol [oral or rectal (suppository)] every 4 hours, or use a prescription medication as directed by your Physician.

### DIET

- 1. Clear liquid diet for first 8 hours (tea, ice chips, apple juice, Jell-O, broth).
- 2. Increase diet gradually avoid greasy, spicy foods during the first 24 hours.

### **ACTIVITY**

1. No heavy lifting or bending until your 1st post-op appointment.

## EXPECTATIONS DURING THE FIRST THREE WEEKS

- 1. Leave patch on until directed by your Physician.
- 2. After patching has been discontinued, apply the topical antibiotic as directed. Wash your hands before and after applying the medication. A mild discharge is not unusual at this stage.
- 3. There is a conformer placed behind the eyelids to maintain the space needed for the prosthesis (false eye). It will appear clear or translucent, and you may see drainage holes in the material.
- 4. After sufficient healing (about six weeks) a prosthesis will be fitted to match the appearance of the other eye.

Weekdays from 8:30 a.m. to 5 p.m. call the Eye Department at 571-4022, or outside normal hours, call the Hospital Operator at 571-4000 to speak to the ophthalmologist on call if you have:

- 1. Increasing pain, swelling, or discharge
- 2. Fever

### FOLLOW UPAPPOINTMENT

Please keep your follow-up appointment located in your Pre-op Packet.

The instructions on this form have been explained to me. I understand them and have received a copy of them.

SIGNATURE OF PATIENT OR LEGAL REPRESENTATIVE	DATE