

Pediatric Well Visit and Vaccine Schedules

Age	Age Range	Vaccine Schedule	Laboratory Evaluation
2-3d		no vaccines	
2w	Before 3 weeks of age	no vaccines	
2m	Before 2 1/2 mos. of age	Pediarix or Dtap, IPV, HEP B and Hib, Prevnar, Rotavirus	
4m	Before 5 mos. of age	Pediarix or Dtap, IPV, HEP B and Hib, Prevnar, Rotavirus	
6m	Before 7 mos. of age	Pediarix or Dtap, IPV, HEP B and Hib, Prevnar, Rotavirus, Influenza if appropriate	
9m	Before 10 mos. of age	Review Immunizations per recommended schedule, if UTD no immunizations needed, consider PPD, Influenza if appropriate	
12m	Must be after 1st birthday	MMR, Varicella, HepA, Influenza if appropriate	Hct, screen for risk - lead
15-18m	Before 18 mos. of age	Dtap, Hib, Prevnar, Influenza if appropriate	
22-24m	Before 24 mos. of age	Hep A#2, Influenza if appropriate	
3.5y*	Before 3yrs., 9 mos. of age	Catch up time for immunizations, Prevnar 13 up to 60 months of age if indicated, Influenza if appropriate	
5y	4-6 years of age (must have immun. before kindergarten)	Kinrix or Dtap, IPV and MMR, Varicella, Influenza if appropriate	screen for risk - H/H, lead
7y	6 1/2 - 8 1/2 years of age	Review Immunizations per recommended schedule, if UTD no immunizations needed, Influenza if appropriate	
9y	8 1/2 - 11 years of age	Review Immunizations per recommended schedule, if UTD no immunizations needed, Influenza if appropriate	
11y	Must be after the 11th birthday	Tdap, HPV, Menactra, Influenza if appropriate	
13y	12 - 14 1/2 years of age	Review Immunizations per recommended schedule, if UTD no immunizations needed. Influenza if appropriate	screen females for risk of anemia <i>Urine STD testing on all teens</i>
15y	14 1/2 - 16 1/2 years of age	Review Immunizations per recommended schedule, if UTD no immunizations needed. Influenza if appropriate	<i>Urine STD testing on all teens</i>
17y	16 1/2 - 18 years of age	Menactra booster #2, Influenza if appropriate **	<i>Urine STD testing on all teens</i>

** 1. Give menactra at age 11-12 yrs with booster at 16 yrs. 2. Give 1 dose at 13-18 yrs if not previously vaccinated. 3. If received 1st dose between 13-15 yrs will need booster at 16-18 yrs. 4. 1 dose to any unvaccinated college freshman.

**1. Give Hep B at birth. 2. Give booster 2 month, 4 month and 6 month or Give Hep B at 2 month, 4 month and 6 month.

**1. Give HPV at age 11 years. 2. Give booster at least 2 months after the 1st dose. 3. Give booster at least 4 months after 2nd dose.