

BEFORE SURGERY

Diet Instructions Checklist
- ADULT & CHILD

FAILURE TO FOLLOW THESE INSTRUCTIONS MAY RESULT IN A DELAY OR CANCELLATION OF YOUR SURGERY

IF YOU ARE A DIABETIC, DRINK WATER ONLY

Adult Diet Instructions	Child Diet Instructions
□ At 2 hours before your check-in time, drink 12 ounces of a carbohydrate rich drink. Finish drinking it within 15 minutes. Drink any of the following carbohydrate rich drinks: ~ Gatorade ~ Apple Juice (no pulp)	You may give your child formula or milk, up until 6 hours before check-in time.
	If breastfeeding, you may give your infant breast milk up until 4 hours before check-in time.
	☐ Give your child clear liquids2 hours before check-in time.
 □ DO NOT eat any solid foods after 11PM. However, you may drink the following clear liquids up until 2 hours before your check-in time: ~ Gatorade ~ Apple Juice (no pulp) ~ Water 	 Your child can drink the following clear liquids only: ~ Gatorade ~ Apple Juice (no pulp) ~ Pedialyte ~ Water
	□ DO NOT allow your child to eat any solid foods after 11PM.
☐ DO NOT chew gum, eat candy, or drink tea or coffee after 11PM.	□ DO NOT allow your child to chew gum or eat candy/gummies after 11PM.
☐ STOP drinking alcohol 2 days before surgery.	